

Double cellular activity to firm the skin

305% Synthesis of Laminin-5

19% Increased cutaneous density

405% Expression of α 6-Integrin



Description

Hexapeptide from an adhesion sequence of laminin, which restores the cutaneous structure.

Appearance

Transparent solution containing 0.05% active ingredient.

INCI

Water (Aqua), Hexapeptide-10
Please contact us for information on the preservative system.

Science

Over the years the adhesion between the epidermal and dermal cells is reduced, which results in the loss of skin elasticity. BE Superlift is a firming active that promotes a double-cellular activity on both keratinocytes and fibroblasts. It increases the synthesis of laminin-5, α 6-integrin and hemidesmosomes, thus strengthening the dermo-epidermal junction.

Properties

Redensifying and restructuring effect.

Applications

BE Superlift^{**} can be incorporated in cosmetic formulations such as emulsions, gels, serums where restructuring and refirming of skin is desired.







& Superlift[™]

• Stimulation of adhesion proteins production

Efficacy demonstrated on 3D skin models which were treated with BE Superlift powder 0.005%





 Stimulation o f hemidesmosome formation

The yellow areas correspond to hemidesmosomes.





• Complementary Studies:

Keratinocyte proliferation 37.6% Fibroblast proliferation 74.6% Significant increase of α 6-Integrin and Laminin-5 production

Increment of hemidesmosome number

BE Superlift[™] powder at 0.005% proved to increase the number of hemidesmosomes, thus increasing cohesion between dermis and epidermis.

 \bullet Stimulation of α 6-Integrin and Laminin-5 pr oduction

Laminin-5 and α 6-Integrin expression was determined by immunohistochemistry Skin biopsies of three patients were carried out before and after a two-month treatment with a cream containing 10% BE Superlift Solution.









0 MONTHS



Green immunofluorescent stain for α 6-Integrin


305% Synthesis of Laminin-5 405% Expression of α6-Integrin

Evaluation of skin appear ance

Panel of 20 female volunteers

A cream containing 10% BE Superlift[™] Solution was applied twice a day during 60 days.


A dermatologist evaluated several parameters.




Evaluation of the dermis density

Panel of 20 female volunteers aged 55 to 64.

A placebo cream was applied twice daily on one side of the face and the cream containing 10% BE Superlift Solution on the other, during 54 days.



BE Superlift[™] solution 10%



BE Superlift solution 10%

- Skin appearance is significantly increased after 60 days
 - +40% Skin compactness
 - +36% Skin tonicity
 - +45% Skin smoothness



+19% Incr eased cutaneous density after 54 days